

insect answers



LILAC LEAFMINER

The lilac leafminer, *Gracillaria syringella*, (leaf blotch miners, family Gracillariidae) attacks nearly all varieties of lilac. It is also known to attack privet, ash, deutzia, and euonymus. The moth originated in Europe and inhabits eastern Canada, northeastern United States, and the Pacific Northwest.

Life History

Adult moths emerge in the spring from overwintering pupae in old leaves and soil. They are small with a wingspan of 2/5 inch (10 mm). The wings are brown with six irregular yellow lines. The female deposits small groups of eggs along the midrib and other veins on the undersides of leaves. Upon hatching, the new larva enters the leaf directly under the egg shell and creates a linear mine which cannot be seen from the top of the leaf. The second larval stage begins to form the unsightly blotch mines. Later the larvae leave the mines and roll the leaves. The larvae are glossy green, and there may be several larvae per mine. When the larval population is especially heavy, the brown, unsightly leaves will drop prematurely. When mature, the larvae drop to the soil on silken threads and pupate. In midsummer, the moths emerge and lay eggs for a second generation which overwinter as larvae or pupae. There may be a third generation in late summer, in which case this would be the overwintering generation.

Control

If only an occasional leaf is infested, pick it off and destroy it. Since the larvae pupate



Lilac leaves infested with lilac leafminer larvae.

among litter and old leaves, future problems can be avoided by clean garden practices. Rake up and destroy old leaves. Chemical control may be necessary if unsightliness becomes intolerable (or the plant is suffering serious defoliation).

Diazinon, malathion, or Orthene are recommended for control of this pest. Spray when blotches are first noticed or at least before the larvae roll the leaves. Repeat applications may be necessary. Be sure the plant or a category which includes the plant is listed on the label of the insecticide you choose. The commonly available malathion products do not have lilac leafminer listed, but they may be used provided the plant is indicated.

By S. J. Collman, WSU Cooperative Extension Agent, King County, and A. L. Antonelli, Extension Entomologist, WSU Western Washington Research and Extension Center, Puyallup. Issued by Washington State University Cooperative Extension, J. O. Young, Director, and the U.S. Department of Agriculture, in furtherance of the Acts of May 8 and June 30, 1914. Cooperative Extension programs and employment are available to all without discrimination. Trade names have been used to simplify the presentation of information. No endorsement is intended. Published February 1982.



